

Generalized Instantaneous Bethe–Salpeter Equation and Exact Quark Propagators

Wolfgang Lucha* and Franz F. Schöberl†

*Institute for High Energy Physics, Austrian Academy of Sciences,
Nikolsdorfergasse 18, A-1050 Vienna, Austria

E-mail: wolfgang.lucha@oeaw.ac.at

†Department of Theoretical Physics, University of Vienna,
Boltzmanngasse 5, A-1090 Vienna, Austria

E-mail: franz.schoeberl@univie.ac.at

Assuming all interactions to be instantaneous *and* the exact fermion propagator S to be approximately given by $iS^{-1}(p) = A(\mathbf{p}^2)\not{p} - B(\mathbf{p}^2)$, i.e., with its scalar functions A and B depending only on \mathbf{p} , a three-dimensional reduction of the homogeneous Bethe–Salpeter equation retaining, in contrast to the Salpeter equation, the exact propagators (crucial for, e.g., a proper incorporation of dynamical chiral symmetry breakdown) is constructed [1]. For spherically symmetric interactions the resulting bound-state equation reduces to a set of (coupled) integral equations for the independent radial components of the bound-state amplitude [2], which may be solved by conversion into an equivalent matrix problem [3]. Adopting as light-quark propagator the solution of the quark Dyson–Schwinger equation found within a model [4] consistent with the QCD axial-vector Ward–Takahashi identity, a tentative application [5] to “pion-like” pseudoscalar ($J^{PC} = 0^{-+}$) mesons considered as quark–antiquark bound states formed by linear confining interactions of time-component Lorentz-vector Dirac structure $\gamma^0 \otimes \gamma^0$ indicates that our “exact-propagator instantaneous formalism” yields significantly smaller spacings of its bound-state mass eigenvalues than those obtained from Salpeter’s equation for “reasonable” constituent light-quark masses.

REFERENCES

1. W. Lucha and F. F. Schöberl, *J. Phys. G: Nucl. Part. Phys.* **31**, 1133 (2005) [hep-th/0507281].
2. J.-F. Lagaë, *Phys. Rev. D* **45**, 305 (1992); M. G. Olsson, S. Veseli, and K. Williams, *Phys. Rev. D* **52**, 5141 (1995) [hep-ph/9503477].
3. W. Lucha, K. Maung Maung, and F. F. Schöberl, *Phys. Rev. D* **63**, 056002 (2001) [hep-ph/0009185]; “Instantaneous Bethe–Salpeter Equation: (Semi-)Analytical Solution,” in *Proceedings of the Int. Conf. on “Quark Confinement and the Hadron Spectrum IV,”* edited by W. Lucha and K. Maung Maung, World Scientific, Singapore, 2002, pp. 340 [hep-ph/0010078]; *Phys. Rev. D* **64**, 036007 (2001) [hep-ph/0011235]; W. Lucha and F. F. Schöberl, *Int. J. Mod. Phys. A* **17**, 2233 (2002) [hep-ph/0109165].
4. P. Maris and C. D. Roberts, *Phys. Rev. C* **56**, 3369 (1997) [nucl-th/9708029]; P. Maris and P. C. Tandy, *Phys. Rev. C* **60**, 055214 (1999) [nucl-th/9905056]; P. Maris, “Continuum QCD and Light Mesons,” in *Proceedings of the Int. Conference on “Quark Confinement and the Hadron Spectrum IV,”* edited by W. Lucha and K. Maung Maung, World Scientific, Singapore, 2002, pp. 163 [nucl-th/0009064].
5. Li Z.-F., W. Lucha, and F. F. Schöberl, *Mod. Phys. Lett. A* **21**, 1657 (2006) [hep-ph/0510372].